

SUTAB®

(sodium sulfate, magnesium sulfate, and potassium chloride)

Tablets

1.479 g/0.225 g/0.188 g

DEPOSIT DAY OF PROCEDURE: \$ _____

Appointment Date: ___/___/___ Time: _____ AM/PM Arrival Time: _____ Location: _____

NOTE: Before beginning the following steps, **have your prescription for filled at the pharmacy.**

*****Please advise your physician prior to your appointment if you have any problems with your heart valve or if you have a heart murmur.*****

You cannot drive the day of the procedure

5 days before procedure: DO NOT TAKE Aspirin or aspirin type products: Coumadin (Warfarin), Advil, Aleve, Motrin, Ecotrin, Plavix (Clopidogrel bisulfate), Xarelto (rivaroxaban), Effient (Prasugrel), Pradaxa (dabigatran), Eliquis (apixaban). **You will need clearance from your Cardiologist prior to stopping these medications.**

DO NOT EAT food containing seeds for **five days prior** to your procedure which included but not limited to sesame, poppy seeds, strawberries, pepper, etc.

*******Any medication for diabetes should be brought along with you the day of the procedure to be taken after the procedure is completed- Check with your physician about possibly adjusting you dosage prior to procedure******

THE DAY BEFORE YOUR PROCEDURE

Breakfast: A low residue diet (Eggs or plain yogurt). Clear Liquids the rest of the day: Soup broth, Jello, Gatorade
NOTHING RED IN COLOR.

3 PM Dose 1

Step 1: Open 1 Bottle of 12 tablets

Step 2: Fill container up to fill line.

Step 3: Swallow each tablet with a sip of water, and drink the entire amount of water within 15-20 minutes.

Step 4: Approx 1 hr after last tablet fill container and drink within 30 minutes.

Step 5: Approx 30 minutes after finishing fill again and drink entire container within 30 minutes.

**** If nausea, excessive bloating or cramping begins, slow down drinking the additional water.**

11 PM Dose 2

Step 1: Open 1 Bottle of 12 tablets

Step 2: Fill container up to fill line.

Step 3: Swallow each tablet with a sip of water, and drink the entire amount of water within 15-20 minutes.

Step 4: Approx 1 hr after last tablet fill container and drink within 30 minutes.

Step 5: Approx 30 minutes after finishing fill again and drink entire container within 30 minutes